

META-ANALYSES OF THE EPIDEMIOLOGICAL EVIDENCE RELATING ETS TO LUNG CANCER AND HEART DISEASE

INTRODUCTION

In two separate documents, the epidemiological evidence relating to possible effects of ETS in never smokers is presented, on an individual study basis, from over 60 lung cancer studies and from almost 30 heart disease studies. In attempting to assess this overall evidence, the technique of meta-analysis is often used to combine the relative risk estimates from the individual studies. Results from a number of meta-analyses are presented in this document. However, before presenting these results, attention should be drawn to some features and limitations of the technique.

Meta-analysis was originally designed to combine evidence from randomised studies of similar design, to try to obtain a more accurate estimate of the possible effect of treatment. However, in recent years it has been increasingly used to combine evidence from epidemiological studies of quite widely varying design. It is important to realise that such a procedure may result in a combined relative risk estimate that has narrow confidence limits and therefore appears precise, but is in fact a quite inaccurate estimate of the true effect of treatment (if any).¹

There are three major potential problems with meta-analysis.

The first, which also applies to the meta-analysis of randomised studies, lies in the possibility that the studies being analysed are unrepresentative of all those studies that actually exist. If, for example, editors are more likely to publish studies showing positive results than they are to publish studies showing negative results, and meta-analysis is based only on published evidence, such "publication bias" will lead to over-estimation of the true effect of treatment. Publication bias can result from various other sources also, and though various methods have been proposed for detecting and correcting for it, all have limitations.²

The second problem, specific to epidemiological studies, is that the results of the individual studies may be distorted by confounding and the other sources of bias that would be avoided in randomised studies. Some types of bias may be common to all (or most) of the studies. For example, it would be expected that studies would tend to find a higher risk of liver cirrhosis in heavy smokers than in nonsmokers, simply because heavy smokers are more likely to be heavy drinkers. Meta-analysis would not remove such a bias, and therefore may be misleading. A statistically significant result from a meta-analysis of epidemiological data does not necessarily establish that any underlying effect exists.^{3,4}

The final problem, particularly important in epidemiological studies, arises because of differences between the studies being combined. Subjects classified as exposed may, for example, have higher average exposure to the agent of interest in some studies than others, so that, even in the absence of any bias, estimates of the true effect of exposure might not be expected to be the same in each study. Many other features of the study are also likely to add to the heterogeneity. These include aspects of study quality, such as the type of study design used and the accuracy of measurement of exposure and disease. They also include the nature of the population studied and where and when the study was conducted. If one is calculating a single combined estimate based on heterogeneous data - averaging apples and oranges in a sense - there is an obvious difficulty in interpreting the combined meta-analysis estimate.⁵

There are two main methods of conducting meta-analysis. One approach, **fixed-effects meta-analysis**, is to assume all the individual study estimates derive from a common mean, with their contribution to the overall estimate depending only on within-study variability, with large studies carrying more weight than do small studies. The alternative approach, **random-effects meta-analysis**, assumes that the individual study estimates derive from a distribution of effects, with the weighting of the individual estimates depending both on the within-study and the between-study variability. Both methods have their advantages and disadvantages.⁶ While the fixed-effects method totally ignores the not at all unlikely possibility of systematic heterogeneity between studies, the attempt to take heterogeneity into account by the random-effects method depends on an assumption about the nature of the distribution of effects over studies that is difficult to justify. Where heterogeneity exists, it is often held that more is to be gained by carefully examining its possible sources than by conducting random-effects meta-analysis.^{1,7,8} If, for example, the true situation is that the effect is consistently X in one group of studies and Y in another, then it would seem better to report the results as such than to attempt to produce a combined estimate.

While meta-analysis has its problems, it nevertheless remains of some value. One might regard the resulting estimate as an indicator of the approximate magnitude of the association to be explained, to be compared with estimates of the approximate magnitude of the various biases that might exist. In the context of ETS and lung cancer, for example, Hackshaw *et al*⁹ have claimed that the major sources of bias are insufficient to explain the magnitude of association, a view which has been hotly contested.¹⁰

Combined estimates of relative risk for the various indices of exposure, as estimated by both fixed- and random-effects meta-analysis¹¹ are set out below.

LUNG CANCER

Combining the overall incidence for each of the four most commonly used indices of ETS exposure, the meta-analysis results show no association with childhood or social exposure, but a possible weak association with smoking by the spouse and in the workplace.

Index of ETS exposure	Estimates combined	Meta-analysis relative risk (95% confidence limits)	
		Fixed effects	Random-effects
Workplace exposure	30	1.21 (1.11-1.31)	1.21 (1.11-1.31)
Childhood exposure from any cohabitant	29	1.06 (0.98-1.15)	1.18 (1.00-1.40)*
Childhood exposure from the mother specifically	9	0.96 (0.77-1.20)	0.98 (0.77-1.25)
Social exposure	12	1.04 (0.92-1.17)	1.02 (0.80-1.28)
Smoking by the wife	20	1.19 (0.99-1.43)	1.19 (0.99-1.43)
Smoking by the husband	61	1.18 (1.12-1.26)	1.24 (1.14-1.35)

*Inflated by one study¹² reporting an extremely high estimate of 12.0 (4.30-32.0) for women.

Subdividing the data on smoking by the husband produces the following:

Studies	Estimates combined	Meta-analysis relative risk (95% confidence limits)	
		Fixed effects	Random-effects
Estimates published in 1981-1989	25	1.38 (1.23-1.55)	1.38 (1.23-1.55)
Estimates published in 1990-2002	36	1.12 (1.05-1.20)	1.18 (1.06-1.31)

It is notable that studies published since 1989 show a statistically significantly lower relative risk than those published earlier.

The meta-analysis relative risks for smoking by the spouse and for workplace ETS exposure are statistically significant. However, the associations are weak and may be explained by various sources of bias, including:

- Confounding**
 Many of the studies on ETS and lung cancer fail to consider diet, lifestyle, family medical history, socio-economic status and other factors known to differ between smoking and nonsmoking households.^{13,14}
- Misclassification**
 Some of the subjects classified as lifelong non-smokers will in fact be current or past smokers who have failed to report this.^{15,16}
- Recall bias**
 The studies rely on reported rather than objectively measured ETS exposure data, which may be affected by presence or knowledge of disease.¹⁰

- **Publication bias**

Studies finding a negative relationship of ETS exposure with lung cancer may be less likely to report their findings than those that find a positive relationship.²

When all the results are considered, and even when meta-analysis is applied, claims that the epidemiological data for lung cancer support an inference of causality⁹ cannot be convincingly justified.¹⁰

HEART DISEASE

The technique of meta-analysis has also been used to combine the results of epidemiological studies on ETS and ischaemic heart disease. The breakdown of results in relation to smoking by the spouse and to workplace ETS exposure can be summarised as follows:

Studies	Estimates combined	Meta-analysis relative risk (95% confidence limits)	
		Fixed-effects	Random-effects
Spouse ever smoked*	39	1.09 (1.05-1.12)	1.17 (1.10-1.24)
Spouse current smoker**	39	1.10 (1.07-1.13)	1.19 (1.12-1.27)
Workplace ETS exposure	16	1.10 (1.00-1.22)	1.10 (1.00-1.22)

* Using estimates for “spouse ever smoked” where a study also provides data for “spouse current smoker”.

** Using estimates for “spouse current smoker” where a study also provides data for “spouse ever smoked”.

As for lung cancer, heart disease studies published in recent years show a weaker relationship of risk to smoking by the spouse than previously published studies. It is notable that the relative risks from one very large US study published in 1995 were very close to 1.00 in each sex, and not statistically significant. This study provided data on some 15,000 heart disease cases, similar to the total number in all the other studies combined.

While the overall adjusted relative risk estimates for spousal smoking are statistically significant, they are based on heterogeneous estimates which are substantially higher in small than in large studies. Many of the studies failed to control adequately for confounding or the various other sources of bias present in such epidemiological studies, none adjusting for misclassification of smoking habits. Heart disease studies show no clearly significant relationship with workplace ETS exposure.

Again, claims that the epidemiological data for heart disease support an inference of causality^{17,18} cannot be convincingly justified.¹⁹

ETS META-ANALYSES FOR LUNG CANCER AND HEART DISEASE

Endpoint	Estimates included	Number of estimates	Relative risk (95% confidence limits)	
			Fixed-effects	Random-effects
LUNG CANCER				
Husband's smoking	All	61	1.18 (1.12-1.26)	1.24 (1.14-1.35)
	USA [†]	19	1.13 (1.02-1.25)	1.13 (1.02-1.25)
	Europe	13	1.29 (1.12-1.48)	1.29 (1.12-1.48)
	Asia	29	1.19 (1.09-1.29)	1.30 (1.13-1.49)
	1981-86	12	1.30 (1.11-1.52)	1.30 (1.09-1.55)
	1987-89	13	1.49 (1.26-1.77)	1.49 (1.26-1.77)
	1990-94	11	0.99 (0.89-1.09)	1.04 (0.84-1.28)
	1995-98	11	1.23 (1.08-1.40)	1.25 (1.07-1.47)
	1999-02	14	1.25 (1.10-1.42)	1.25 (1.10-1.42)
Wife's smoking	All	20	1.19 (0.99-1.43)	1.19 (0.99-1.43)
	USA [†]	9	1.14 (0.85-1.53)	1.14 (0.85-1.53)
Husband or wife smoked	All	76	1.19 (1.12-1.25)	1.24 (1.15-1.34)
	USA [†]	26	1.13 (1.03-1.25)	1.13 (1.03-1.25)
	Europe	16	1.27 (1.12-1.43)	1.27 (1.12-1.43)
	Asia	34	1.19 (1.09-1.29)	1.29 (1.13-1.48)
	1981-86	18	1.32 (1.14-1.53)	1.32 (1.13-1.54)
	1987-89	15	1.52 (1.28-1.80)	1.52 (1.28-1.80)
	1990-94	12	0.99 (0.89-1.10)	1.05 (0.85-1.29)
	1995-98	16	1.21 (1.08-1.37)	1.22 (1.07-1.39)
	1999-2002	15	1.22 (1.09-1.38)	1.22 (1.09-1.38)
Workplace ETS exposure	All	30	1.21 (1.11-1.31)	1.21 (1.11-1.31)
Childhood ETS exposure				
	From any cohabitant	All	29	1.06 (0.98-1.15)
From the mother specifically	All	9	0.96 (0.77-1.20)	0.98 (0.77-1.25)
ETS exposure in social settings	All	12	1.04 (0.92-1.17)	1.02 (0.80-1.28)
HEART DISEASE				
* Spouse ever smoked	All	39	1.09 (1.05-1.12)	1.17 (1.10-1.24)
	1984-88	7	1.22 (1.04-1.44)	1.26 (1.02-1.55)
	1989-92	11	1.29 (1.17-1.43)	1.39 (1.18-1.64)
	1993-96	9	1.03 (0.99-1.07)	1.04 (0.99-1.08)
	1997-2002	12	1.20 (1.12-1.27)	1.24 (1.09-1.40)
	1-99 cases	5	1.53 (1.18-2.00)	1.55 (1.17-2.05)
	100-199	10	1.37 (1.11-1.69)	1.37 (1.11-1.69)
	200-999	14	1.28 (1.15-1.43)	1.32 (1.13-1.55)
	1000+	10	1.06 (1.03-1.09)	1.08 (1.02-1.14)
Spouse current smoker	All	39	1.10 (1.07-1.13)	1.19 (1.12-1.27)
Workplace ETS exposure	All	16	1.10 (1.00-1.20)	1.10 (1.00-1.22)

Note: The individual study data used for the meta-analyses are given in the documents "*Epidemiological Evidence on Environmental Tobacco Smoke and Lung Cancer*" and "*Epidemiological Evidence on Environmental Tobacco Smoke and Heart Disease*", or their associated appendices. The individual study estimates used are those adjusted for as many potential confounding variables as the authors have presented results for.

* Studies by Palmer and Mannino omitted as confidence limits not available, and by Spence omitted as neither relative risk nor confidence limit available.

† Includes one study in Canada.

REFERENCES

1. Egger M, Schneider M, Davey Smith G: Spurious precision? Meta-analysis of observational studies. *BMJ* **316**:140-144,1998
2. Thornton A, Lee P: Publication bias in meta-analysis: its causes and consequences. *J Clin Epidemiol* **53**:207-216,2000
3. Shapiro S: Meta analysis/Shmeta-analysis. *Am J Epidemiol* **140**:771-778,1994
4. Petitti DB: Of babies and bathwater. *Am J Epidemiol* **140**:779-782,1994
5. Peto R: Why do we need systematic overviews of randomized trials? [Transcript of an oral presentation, modified by the editors]. *Stat Med* **6**:233-244,1987
6. Jones DR: Meta-analysis: Weighing the evidence. *Stat Med* **14**:137-149,1995
7. Thompson SG: Why sources of heterogeneity in meta-analysis should be investigated. *BMJ* **309**:1351-1355,1994
8. Blair A, Burg J, Foran J, *et al*: Guidelines for application of meta-analysis in environmental epidemiology. *Regul Toxicol Pharmacol* **22**:189-197,1995
9. Hackshaw AK, Law MR, Wald NJ: The accumulated evidence on lung cancer and environmental tobacco smoke. *BMJ* **315**:980-988,1997
10. Lee PN, Fry JS, Forey BA: Revisiting the association between environmental tobacco smoke exposure and lung cancer risk. V. Overall conclusions. *Indoor Built Environ* **11**:59-82,
11. Fleiss JL, Gross AJ: Meta-analysis in epidemiology, with special reference to studies of the association between exposure to environmental tobacco smoke and lung cancer: a critique. *J Clin Epidemiol* **44**:127-139,1991
12. Rapiti E, Jindal SK, Gupta D, *et al*: Passive smoking and lung cancer in Chandigarh, India. *Lung Cancer* **23**:183-189,1999
13. Thornton A, Lee P, Fry J: Differences between smokers, ex-smokers, passive smokers and non-smokers. *J Clin Epidemiol* **47**:1143-1162,1994
14. Fry JS, Lee PN: Revisiting the association between environmental tobacco smoke exposure and lung cancer risk. II. Adjustment for the potential confounding effects of fruit, vegetables, dietary fat and education. *Indoor Built Environ* **10**:20-39,2001
15. Lee PN, Forey BA: Misclassification of smoking habits as determined by cotinine or by repeated self-report - a summary of evidence from 42 studies. *J Smoking-Related Dis* **6**:109-129,1995
16. Lee PN, Forey BA, Fry JS: Revisiting the association between environmental tobacco smoke exposure and lung cancer risk. III. Adjustment for the biasing effect of misclassification of smoking habits. *Indoor Built Environ* **10**:384-398,2001
17. Glantz SA, Parmley WW: Passive smoking and heart disease. Mechanisms and risk. *JAMA* **273**:1047-1053,1995

18. Law MR, Morris JK, Wald NJ: Environmental tobacco smoke exposure and ischaemic heart disease: an evaluation of the evidence. *BMJ* **315**:973-980,1997
19. Lee PN, Roe FJC: Environmental tobacco smoke exposure and heart disease: a critique of the claims of Glantz and Parmley. *Hum Ecol Risk Ass* **5**:171-218,1999